

SUPERLUMINAL INFLUENCES, HIDDEN VARIABLES, AND SIGNALING

Valerio Scarani*, Nicolas Gisin

Group of Applied Physics, University of Geneva

20, rue de l'Ecole-de-Médecine, CH-1211 Geneva, Switzerland

Abstract

We consider alternative models to quantum mechanics, that have been proposed in the recent years in order to explain the EPR correlations between two particles. These models allow in principle local hidden variables produced at the source, and some superluminal "hidden communication" (or "influences") to reproduce the non-local correlations. Moving to the case of three particles, we show that these alternative models lead to signaling when "hidden communication" alone is considered as the origin of the correlations.

1 Introduction

In the classical world, we are accustomed to two kinds of causes for correlations: (i) A manufacturer produces white and black balls; he sells them by pairs, and he always puts a white and a black ball in each box. Here, the source of the perfect anticorrelation is pre-established in the fabric. (ii) A referee decides that the match is over, and sends a signal (whistle) to all players, who stop running. The correlated behavior of the players is the result of having received a signal. However, neither of these models can explain the *Einstein-Podolski-Rosen (EPR) correlations*, that are, the correlations between two quantum entangled particles. In fact, on the one hand, these correlations are established for space-like separated events, ruling out the explanation through the reception of a common signal [1, 2, 3]; on the other hand, their explicit dependence on the meaningful parameters of the experiments cannot be the consequence of sharing a common information pre-established at the source — this is Bell's theorem [4, 5].

The physics community is accustomed since more than one century to the counter-intuitive character of some predictions of quantum mechanics (QM). Still, facing EPR correlations, some physicists are not satisfied: as John Bell put it, "correlations cry out for explanation", and not only for "description". That's why in the last decades some models have been proposed that provide a dynamical explanation to quantum correlations between two particles [6, 7, 8, 9, 10, 11]. These models incorporate Bell's theorem: the correlations are not entirely pre-established at the source; but they admit the existence of some form of communication between the particles, that we shall call here *hidden communication*. This communication must propagate at a speed

*corresponding author. Tel. +41 22 7026883; fax +41 22 7810980; e-mail: valerio.scarani@physics.unige.ch

v_{hc} which is superluminal, to be consistent with experiments performed in the configuration of space-like separation. But one cannot define a superluminal communication in a relativistic space-time [12], without allowing the causal loop or backward-in-time signaling. Therefore the model must single out the frame in which the hidden communication occurs.

Among these dynamical models, some are "alternative models", since the explanation that is proposed leads also to predictions that *differ* in some cases from those of QM, without contradicting the experimental data that are available at the time of the proposal. These models are worth of consideration: they lead to design new experiments, that can discriminate between them and standard QM. If the experiments falsifies QM, a door is opened for new physics; but even if QM is confirmed, which is probable *a priori*, the investigation of the alternative model will have shed new light on the intimate connection between QM and relativity.

We mentioned the astonishing features of EPR correlations, but there is another element that prevents these correlations to become something too dramatic. In spite of their "non-local" character, EPR correlations do not allow *signaling*, that is, sending a message at an arbitrary high speed. More precisely, these correlations alone cannot be used to send a signal at any speed; and classical communication must be established between the two observers if they want to check their correlations. Now, any alternative model, in addition to reproduce QM for all the experiments performed to date, is also supposed to fulfill the no-signaling condition. A model that fails to comply with this requirement would be in open contradiction with special relativity, thus acquiring a very problematic status — although conclusive falsification can come only through experiment.

In this paper, we consider alternative models built on the idea of superluminal hidden communication. We show that if correlations are built only by this communication, these models imply signaling. Thus, any alternative model must invoke both superluminal hidden communication and local variables.

We begin with some general considerations about the no-signaling condition, and the constraints that it imposes.

2 From two to three particles

Consider two quantum particles of arbitrary spin prepared in an entangled state, then flying apart from one another, each to one observer, Alice or Bob. Each observer makes a measurement. Let's call a_j , $j = 1, \dots, J$ the possible outcomes for Alice's measurement; and b_k , $k = 1, \dots, K$ the possible outcomes for Bob's measurement. The joint probability for two outcomes is written $P(a_j, b_k)$; the marginal distributions on Alice's and Bob's side are $P(a_j) = \sum_k P(a_j, b_k)$, respectively $P(b_k) = \sum_j P(a_j, b_k)$.

Now, Alice could decide to make another measurement, whose outcomes are labelled a'_l , with $l = 1, \dots, L$. The no-signaling condition here means that $P'(b_k) = \sum_l P(a'_l, b_k)$ must be equal to $P(b_k)$: on his own side, Bob cannot notice that Alice has chosen another measurement. Of course, the symmetric condition must yield on Alice's side: Alice's marginal distribution cannot be modified by Bob's choosing another measurement.

Now, within the framework of QM, "another measurement" means to measure another observable, or more generally to perform a different POVM. But here we are willing to consider alternative models to QM,




Figure 1: Three different probability distributions for the outcomes of Alice and Bob ($a_j = 0, 1; b_k = 0, 1, 2$), that satisfy the no-signaling condition. The probability of each outcome is measured by the area of the corresponding sector of the unit square. The marginal distributions $P(a_j)$ and $P(b_k)$ are the same in the three cases, while the correlations are completely different; (b) describes the absence of correlation.

that give different predictions for the EPR correlations. In this case, "another measurement" may mean that Alice modifies her parameters so that the correlations are no longer those predicted by QM, that is $P(a_j, b_k) \neq P^{QM}(a_j, b_k)$. Therefore, in the more general framework in which we work, the no-signaling condition implies also that $P(a_j) = P^{QM}(a_j)$, and $P(b_k) = P^{QM}(b_k)$.

Now, in the case of two particles, a wide range of alternative models can be made compatible with the no-signaling condition. This can be easily seen by looking at a graphic representation. In fig. 1 we represent three possible probability distributions for the joint outcomes of Alice and Bob; a probability is measured by the area of the corresponding sector of the unit square. We see that very different situations can be met while conserving the same marginal distributions, that is, without violating the no-signaling condition. In particular, a model that predicts a *complete loss of correlations* is possible: if $P(a_j, b_k) = P^{QM}(a_j)P^{QM}(b_k)$ for all measurements, clearly the marginal distribution on each side is independent on what is done on the other side, and agrees with the prediction of QM.

Thus, as long as one considers only correlation between two particles, the no-signaling condition puts practically no constraint on the correlations. However, two-particle correlations can be seen as marginal distributions of a three-particle probability distribution. This will be discussed in detail in the next sections, but we want to give the reader the feeling of what happens. We have now a probability distribution $P(a_j, b_k, c_l)$. The no-signaling condition forces $P(a_j)$, $P(b_k)$ and $P(c_l)$ to agree with the corresponding prediction of QM, just as above. But in addition, the no-signaling condition may also force some *correlations* to agree with those predicted by QM. Specifically, in the next section we shall build, for two alternative models, a situation in which both $P_{AC}(a_j, c_l)$ and $P_{BC}(b_k, c_l)$ must be identical to those predicted by QM, while $P_{AB}(a_j, b_k)$ is free *a priori*. Consider then the simple case of a dichotomic measurement on each of the three particles, for which QM predicts perfectly correlated outcomes: $P^{QM}(000) = P^{QM}(111) = \frac{1}{2}$. For such a measurement, obviously no freedom is left on $P_{AB}(a_j, b_k)$ if all the other marginals must agree with QM: no alternative model can predict anything else than $P_{AB}(00) = P_{AB}(11) = \frac{1}{2}$, $P_{AB}(01) = P_{AB}(10) = 0$.




Figure 2: The events considered for the argument in Model 1 (PF and $v_{hc} < \infty$). Events D are detections. The black lines are light cones, the dotted lines are the cones of the superluminal "hidden-communication". Bob can calculate the correlations B-C in P_{BC} , before he can calculate the correlation A-B: consequently the correlation B-C cannot depend on whether Alice chose to put or not the time delay τ on her detection. The argument can easily be made symmetric.

3 Predictions of two alternative models

3.1 Model 1: preferred frame and finite velocity

The first model that we consider is a natural modification of Bohmian mechanics. In Bohmian mechanics, the outcomes of a measurement are determined by local parameters (the local settings of the measurement and the quantum potential in which the particle propagates), and by a superluminal hidden communication between the correlated particles [6]. The hidden communication takes place in a preferred frame (PF), an assumption that can be made for quantum phenomena without contradicting relativity in the macroworld [13]. If the speed of this hidden communication in the PF is $v_{hc} = \infty$, the predictions of Bohmian mechanics reproduce perfectly those of QM. However, in the same line of thought one may suppose that $v_{hc} < \infty$. In this case, the model gives some predictions that differ from the QM ones. In fact, the hidden communication sent by the first particle that is detected may not reach the second particle before its own detection. In this case, non-local correlations should disappear. This defines the model.

As we demonstrated in section 2, a complete loss of correlations can be made consistent with the no-signaling condition for two entangled particles, that is, for the original EPR-Bohm experiment. Eberhard [14] was the first to explore the predictions of such a model in the case of three particles: he noticed that signaling could be allowed, at least in the particular version of the model that he built.

We also consider a prediction of this model for an experiment involving three particles. The three entangled particles are labelled A, B and C, and we name Alice, Bob and Charlie the experimenters that perform the measurements. The event "detection of particle X" is written D_X . In the space-time coordinates of the PF, we note

$$D_A = (-x, t_A), \quad D_B = (x, t_B), \quad D_C = (0, t_C). \quad (1)$$

If the experimenters know which frame is the PF, they can arrange the timings of the detection events in such a way that: (I) D_A and D_B are simultaneous: $t_A = t_B = 0$; (II) particle C is detected later, and we suppose that the following condition holds:

$$\frac{3x}{v_{hc}} < t_C < \frac{x}{c}. \quad (2)$$

The upper bound simply means that no classical signal sent by A or B at $t = 0$ might have reached C at t_C ; the lower bound is justified below. Condition (2) is not contradictory as long as $v_{hc} > 3c$, which cannot be taken as absolutely granted since we don't know the PF. However, for two reasonable candidates of PF (the laboratory frame, and the frame of the cosmic background radiation), recent experiments provided a lower bound for v_{hc} which is several orders of magnitude higher than c [15]. So let's assume that $v_{hc} > 3c$ [16]. Condition (2) implies that particle C has received the hidden information from both A and B, since $t_C > \frac{x}{v_{hc}}$, but no classical information. We claim that the correlations A-C and B-C must be those predicted by QM, in order to avoid signaling.

To prove this claim, we refer to fig. 2, in which the argument is sketched for the correlation B-C. $E(r_B r_C)$ can be calculated by Bob after the space-time point $P_{BC} = (x, t_C + \frac{x}{c})$. At that time, Bob cannot have received any classical information from Alice, since an information propagating from D_A to D_B at the speed of light cannot arrive before $P_{AB} = (x, \frac{2x}{c})$. Therefore, the correlation $E(r_B r_C)$ cannot change whatever Alice does, otherwise we would allow signaling from Alice to the pair Bob-Charlie.

Suppose now that Alice delays the detection of particle A by a time $\tau = \frac{x}{v_{hc}} + \epsilon$, so that the event "detection of A" is now D'_A . In this case, A receives the hidden information from B, and C still receives from both — this is the justification of the lower bound in (2). So now we have a normal time ordering $B \rightarrow A \rightarrow C$, thus QM should apply. Consequently, the only way to avoid signaling is to force correlation B-C to be the one predicted by QM, independently of whether A has or has not received the hidden information from B. A completely symmetrical argument, allowing Bob to delay his detection to D'_B , leads to the conclusion that the correlation A-C must also be the QM one.

Now, what about correlation A-B? Within the model, it depends on the situation: (i) If the detections are simultaneous, i.e. in cases (D_A, D_B) or (D'_A, D'_B) , then A and B are either uncorrelated (if we suppose that correlations arise only through reception of the hidden communication), or at most correlated through local variables; (ii) If one detection is delayed, i.e. in cases (D'_A, D_B) or (D_A, D'_B) , this correlation should agree with QM.

In conclusion: under some circumstances this model predicts the disappearance of non-local correlations between A and B, while A-C and B-C are correlated according to QM.




Figure 3: Experimental setup for the argument in Model 2 (Multisimultaneity). The squares represent the "choice devices". See text for details.

3.2 Model 2: Multisimultaneity

This model was proposed some years ago by Suarez and one of us [11], and developed by Suarez in subsequent papers, where he called it "Multisimultaneity" [17]. Here there is not a unique PF. Rather, there are several meaningful frames: for each particle, the meaningful frame is the rest frame of the massive *choice-device* that it meets. The choice-device may be the detector — in which case the "choice" is a form of "collapse" — or the beam-splitter, again in a Bohm-like interpretation where the particle is really localized in a path, its "choices" being determined by a pilot wave or quantum potential. Experiments have been performed with detectors [18] and with beam-splitters in motion [19], both vindicating QM. Our argument is independent of what a choice-device is.

For two particles, Multisimultaneity works as follows: when a particle meets a choice-device, it considers whether *in the rest frame of this device* the other particle has already met its own device [20]. Thus, according to the state of motion of the choice-devices, we can arrange different timings: (I) *Before-before timing*, in which each particle arrives to its own choice-device, in the rest frame of this device, before the other one. In this case, each particle chooses its outcomes taking into account only local settings: two-particle correlations should disappear completely; (II) *Before-after timing*, in which the "before" particle chooses randomly its output, while the "after" one takes into account the measurement that has been performed on the first one. In the before-after timing, Multisimultaneity reproduces the QM results; and this is the situation achieved in all standard EPR experiments [1, 2, 3], that supported QM. (III) *After-after timing*, which is a rather troubling feature of this kind of theories, and is not easy to define while avoiding the causal loop [17, 18].

Again, let's turn to three particles. Consider the experimental setup sketched in fig. 3. Three particles issued from the source S reach the respective choice-devices. The choice-devices of A and B can be put into opposite motion with speed v , high enough to achieve a before-before configuration [11]. Therefore, when

both devices are at a relative rest, QM should apply; when they are in relative motion, particles A and B have to choose in a before-before configuration, therefore their non-local correlation should disappear. As for particle C: its choice-device is motionless in the laboratory, and the projection of the speed v on the lines A-C and B-C (small black arrow in the figure) is too small to change the timing. To be concrete, we suppose that when C meets its choice-device, A and B have already made their choice in the laboratory frame.

An argument analogous to the one discussed for Model 1 applies here: correlations A-C and B-C must be in any case those predicted by QM. In fact, suppose that Bob's device is at rest: Alice may put her device into motion, and the pair Bob-Charlie should see no difference. Symmetrically, if Alice's device is at rest, Bob should be allowed to put his device into motion without signaling to Alice-Charlie. Correlation A-B is the QM one when both Alice's and Bob's devices are at rest in the laboratory frame; but when at least one device moves, the non-local correlations should disappear. Thus, Multisimultaneity also predicts that in some cases A-B can be at most correlated through local variables, while A-C and B-C are correlated according to QM.

4 Hidden communication alone implies signaling

In the context of alternative models invoking superluminal hidden communication, the content of section 2 can be rephrased as follows. In the case of two particles, it is possible to imagine a model in which the correlations are due only to the hidden communication: if the communication is received, the outcomes of the measurements becomes correlated; if the communication is not received, the outcomes are uncorrelated. Here we show that this is no longer true when we go to three particles.

To simplify the analysis, we assume that on each of the three particles a local dichotomic measurement is performed. Alice's possible results r_A are labelled $\xi_A = \pm 1$, and similarly for Bob's and Charlie's measurements. With this labelling, any probability distribution can be written

$$p(r_A = \xi_A, r_B = \xi_B, r_C = \xi_C) = \frac{1}{8} \left[1 + \xi_A E(r_A) + \xi_B E(r_B) + \xi_C E(r_C) + \xi_A \xi_B E(r_A r_B) + \right. \\ \left. + \xi_A \xi_C E(r_A r_C) + \xi_B \xi_C E(r_B r_C) + \xi_A \xi_B \xi_C E(r_A r_B r_C) \right], \quad (3)$$

where $E(\cdot)$ is the expectation value of the random variable. We have seen that both Models 1 and 2 predict a configuration in which:

Condition 1: $E(r_A)$, $E(r_B)$, $E(r_C)$, $E(r_A r_C)$ and $E(r_B r_C)$ must be those predicted by QM; this is the no-signaling condition in the configurations that we have built.

Moreover, we are supposing that the correlations arise only through reception of the hidden communication, Therefore we add to Models 1 and 2 the following:

Condition 2: $E(r_A r_B) = E(r_A)E(r_B)$: complete loss of correlation.

We provide now examples of quantum states and measurements for which Conditions 1 and 2 lead to negative probabilities. Since we consider dichotomic observables, the meaningful degrees of freedom of each particle can be mapped onto a two-dimensional quantum system (qubit).

First example: This is the example that we have already discussed in a semi-qualitative way in section 2. Take

the GHZ state $\frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)$ [21], with $|0\rangle$ and $|1\rangle$ the eigenstates of σ_z . If we measure σ_z on all particles, the quantum statistics give $E^{QM}(r_A) = E^{QM}(r_B) = E^{QM}(r_C) = 0$, $E^{QM}(r_{ArC}) = E^{QM}(r_{BrC}) = 1$. Condition 2 gives consequently $E(r_{ArB}) = 0$. If we put this into (3), we find in particular $p(++-) = \frac{1}{8}[-1 - E(r_{ArBrC})]$, $p(--+) = \frac{1}{8}[-1 + E(r_{ArBrC})]$. These numbers cannot be both non-negative: there is no three-particle probability distribution that is compatible with both Conditions 1 and 2. Actually for this particular case the only way to have non-negative probabilities is to set $E(r_{ArB}) = 1$ and $E(r_{ArBrC}) = 0$, exactly the values predicted by QM. Thus this example shows that all possible models that would predict a reduction of visibility for all measurements, that is $E(r_{ArB}) = V E^{QM}(r_{ArB})$ with $0 \leq V < 1$, lead to signaling — we are not aware of any explicit model of this kind.

Second example: Take the state $|W\rangle = \frac{1}{\sqrt{3}}(|001\rangle + |010\rangle + |100\rangle)$ [22]. If we measure σ_x on all particles, the quantum statistics give $E^{QM}(r_A) = E^{QM}(r_B) = E^{QM}(r_C) = 0$, $E^{QM}(r_{ArC}) = E^{QM}(r_{BrC}) = \frac{2}{3}$. Condition 2 gives consequently $E(r_{ArB}) = 0$. If we put this into (3), we find in particular $p(++-) = \frac{1}{8}[-\frac{1}{3} - E(r_{ArBrC})]$, $p(--+) = \frac{1}{8}[-\frac{1}{3} + E(r_{ArBrC})]$. Again, these numbers cannot be both non-negative. However here we could recover non-negative probabilities by having $E(r_{ArB}) = \frac{1}{3}$, which is lower than the QM value $E(r_{ArB}) = \frac{2}{3}$, but still different from 0.

The appearance of negative probabilities means that the operation "suppression of the correlation between some partners" is not a completely positive (CP) map. It has been recently shown [23] that if a system allows a description agreeing with QM at a given time, the no-signaling condition forces the dynamics to be described by a CP map.

5 Consequences for alternative models

In the introduction we explained that, in the classical world, correlations seem to be always due either to preparation or to communication. Bell's theorem ruled out the first of these origins for EPR correlations. In this paper, we have ruled out also the hypothesis of a hidden communication as unique cause of the correlations, at least in the most natural alternative models.

One can still hope to develop an alternative model that introduces *both* local hidden variables and hidden communication; we call these *mixed models* [24]. In a mixed model, correlations are established by hidden communication if this communication is allowed to reach the other particles, or using a pre-established information if for some reason the hidden communication has not been received. We explain here the relation between our approach and signaling in mixed models.

Consider again Example 1 of the previous section. For that measurement, only the correlations predicted by QM are compatible with the no-signaling condition. Unfortunately, no hidden communication at all is needed to establish such correlations. Even more generally: all the two-particle correlations predicted by QM for a GHZ state can be described by classical mixtures, since the two-particle partial trace on $|GHZ\rangle$ is $\rho = \frac{1}{2}|00\rangle\langle 00| + \frac{1}{2}|11\rangle\langle 11|$. Consider then the following mixed model: (i) the source produces an equal-weight mixture of the possibilities $|000\rangle$ and $|111\rangle$. (ii) If the hidden communication is allowed to reach the particles, the remarkable three-particle GHZ correlations [21] are established. (iii) If the hidden communication does

not reach the particles, their correlations are determined by the preparation. In this last case, the three-particle correlation will disagree with the prediction of QM, but this is not in conflict with the no-signaling condition as long as we have only three particles.

Of course, the question is, whether by considering other quantum states one can show that also mixed models lead to signaling. To do so, one should show that the no-signaling condition can force a violation of a Bell's inequality. Let us sketch the idea of this in the same configuration described above, in which A and B cannot share the hidden communication and are therefore supposed to be correlated only through preparation.

Alice makes on her side N different measurements, and similarly, Bob makes M different measurements; for simplicity, we assume that Charlie makes only one measurement. Thus the three partners share $N \times M$ probability distributions $P(a_j^{(n)}, b_k^{(m)}, c_l)$. The no-signaling constraint says that $P(a_j^{(n)}, c_l)$ and $P(b_k^{(m)}, c_l)$ must be those predicted by QM, whatever the measurements performed by Alice and Bob; and this requirement puts some constraints on the marginals $P(a_j^{(n)}, b_k^{(m)})$; for instance, we have demonstrated above that $P(a_j^{(n)}, b_k^{(m)}) = P(a_j^{(n)})P(b_k^{(m)})$ is impossible. Suppose now that there is a combination of these two-particle probabilities that defines a Bell's inequality which is violated by the QM probabilities $P^{QM}(a_j^{(n)}, b_k^{(m)})$. One can hope that the constraints put by the no-signaling condition are strong enough to force the violation of the same inequality, though possibly a weaker violation. If this is the case, then no suitable preparation can lead to probabilities $P(a_j^{(n)}, b_k^{(m)})$ compatible with the no-signaling constraints: mixed models would also imply signaling.

We have not found such a state and set of measurements. We note that the study of the violation of Bell's inequalities by partial states [25], as well as the study of Bell's inequalities beyond the case of two qubits [26], are presently open fields of research. So it is likely that in the next years we shall dispose of more adequate tools to tackle the robustness of mixed models against signaling.

Acknowledgements

We thank Ph. Eberhard, S. Popescu and A. Suarez for fruitful discussion, and an anonymous referee for valuable comments. We acknowledge partial financial support from the Swiss FNRS.

References

- [1] A. Aspect et al., *Phys. Rev. Lett.* **49** (1982) 91
- [2] W. Tittel et al., *Phys. Rev. Lett.* **81** (1998) 3563
- [3] G. Weihs et al., *Phys. Rev. Lett.* **81** (1998) 5039
- [4] J. S. Bell, *Speakable and Unspeakable in Quantum Mechanics: Collected papers on quantum philosophy* (Cambridge University Press, Cambridge, 1987).
- [5] There have been several attempts of weakening the importance of Bell's result on the mathematical ground, without success until now. Recently, a proposal by K. Hess and W. Philipp [*Proc. Nat. Acad.*

Sci. **98** (2001) 14228] has attracted some attention, but in our opinion it also suffers from a subtle mistake. In fact, if the correlations at the source are setting-independent as they should, the setting-dependent distribution ρ_{ab} can be built only if both settings **a** and **b** are known in each analyzing station. This means instantaneous hidden communication between the stations.

- [6] D. Bohm, B.J. Hilley, *The undivided universe* (Routledge, New York, 1993)
- [7] Ph. Pearle, *Phys. Rev. D*, **13**, 857, 1976
- [8] G. C. Ghirardi, A. Rimini, and T. Weber, *Phys. Rev. D*, **34**, 470 (1986)
- [9] I. Percival, *Quantum State Diffusion*, Cambridge Univ.Press 1998
- [10] N. Gisin, *Helv. Phys. Acta* **62**, 363, 1989.
- [11] A. Suarez, V. Scarani, *Phys. Lett. A* **232** (1997) 9
- [12] H. Reichenbach, *The philosophy of Space and Time* (Dover, New York, 1958).
- [13] F. Reuse, *Ann. Phys.* **154** (1984) 161; P. Caban, J. Rembielinski, *Phys. Rev. A* **59** (1999) 4187
- [14] Ph. Eberhard, *A Realistic Model for Quantum Theory with a Locality Property*, in: W. Schommers (ed.), *Quantum Theory and Pictures of Reality* (Springer, Berlin, 1989)
- [15] V. Scarani, W. Tittel, H. Zbinden, N. Gisin, *Phys. Lett. A* **276** (2000) 1
- [16] A referee pointed out that one can find a different ordering of the events such that the condition $v > 3c$ can be eliminated, at least for the example of the GHZ state. The situation is the following: particle A leaves on the left, particles B_1 and B_2 on the right. These two particles are measured at nearby locations but simultaneously, so that they cannot be correlated by the hidden communication. If both have received the communication from particle A , they must become correlated, because of the explanation at the end of section 2. If they have not received communication from A and if the correlation arise only through communication, then B_1 and B_2 are uncorrelated. Note that this is a different geometry from the one proposed in the main text. Here, the alternatives are "no pair correlated" vs "pairs $A - B_1$ and $A - B_2$ correlated" (and forcing a correlation on $B_1 - B_2$). In the main text, the alternatives are "pairs $A - C$ and $B - C$ correlated" vs. "all three pairs correlated".
- [17] A. Suarez, *Phys. Lett. A* **236** (1997) 383
- [18] H. Zbinden, J. Brendel, N. Gisin, W. Tittel, *Phys. Rev. A* **63** (2001) 022111
- [19] A. Stefanov, H. Zbinden, N. Gisin, A. Suarez, quant-ph/0110117, to appear in *Phys. Rev. Lett.*
- [20] Note that Multisimultaneity predicts differences with QM also in the case $v_{hc} = \infty$. The case of finite v_{hc} has not been studied within this approach.
- [21] D. Greenberger, M. Horne, A. Zeilinger, in M. Kafatos (ed.), *Bell's Theorem, Quantum Theory, and Conceptions of the Universe* (Kluwer, Dordrecht, 1989).

- [22] W. Dür, G. Vidal, I. Cirac, *Phys. Rev. A* **62** (2000) 062314
- [23] C. Simon, V. Bužek, N. Gisin, *Phys. Rev. Lett.* **87** (2001) 170405
- [24] The Bohm-Bub theory has been shown to be a mixed model requiring some non-local communication, and leading to signaling: T. Durt, *Helv. Phys. Acta* **72** (1999) 356
- [25] V. Scarani, N. Gisin, *Phys. Rev. Lett.* **87** (2001) 137901; *Phys. Rev. A* **65** 012311 (2002)
- [26] D. Collins, N. Gisin, N. Linden, S. Massar, S. Popescu, *Phys. Rev. Lett.* **88** (2002) 040404